



# CLINICAL PRACTICE GUIDELINES AND RECOMMENDATIONS FOR ADULT DIABETES CARE 2009

		Frequency	Comments
<b>History, Physical Examination &amp; Laboratory (screening &amp; diagnostic)</b>	Blood Pressure and Weight	Every 3-6 months	Treat to a systolic BP <130 mmHg target; to a diastolic BP < 80 mmHg target.
	Dilated eye exam	Annually	Refer to ophthalmologist or optometrist
	Foot Exam	Every routine visit	Visual exam without shoes or socks; refer as indicated Note: All patients should be screened for distal symmetric polyneuropathy (DPN) at diagnosis and at least annually. Once diagnosis of DPN is established, special foot care is appropriate.
	Comprehensive lower extremity sensory exam	Initial/annually	Teach protective foot behavior if sensation diminished. Refer to podiatrist if indicated
	Dental exam	Every 6 months	Refer to Dentist
	Cardiovascular exam	Annually	Refer to cardiologist for further evaluation as indicated
	Depression assessment	Annually	Assess for depression with Patient Health Questionnaire-9 (PHQ-9); alternatively screen for depression verbally and if positive, use PHQ-9
	Screening to detect Pre-Diabetes ( <i>IFG or Impaired Fasting Glucose; IGT or Impaired Glucose Tolerance</i> )	As indicated/repeat testing should be carried out at 3-year intervals.	Should be considered for: <ul style="list-style-type: none"> <li>● Asymptomatic adults of any age who are overweight or obese and have additional risk factors for diabetes.</li> <li>● In those without a risk factor, testing should begin at age 45.</li> </ul> Note: for this purpose, this recommendation would apply particularly to those with a BMI $\geq 25$ kg/m <sup>2</sup> if they have another risk factor for diabetes, e.g., habitually physically inactive, have first-degree relative with diabetes, are members of high-risk ethnic population, have delivered a baby weighing >9 lbs or have been diagnosed with gestational diabetes, are hypertensive, have HDL cholesterol level <35 mg/dl (0.90 mmol/l) and/or a triglyceride level >250 mg/dl (2.82 mmol/l), have a clinical condition associated with insulin resistance, or have history of vascular disease.
	<i>Screening to Detect type 2 Diabetes in Children</i>	As indicated, starting at age 10 or at onset of puberty, if puberty occurs at a younger age; should be repeated every 3 years.	Should be considered for: <ul style="list-style-type: none"> <li>● Overweight (BMI &gt;85<sup>th</sup> percentile for age and sex, weight for height &gt;85<sup>th</sup> percentile, or weight &gt;120% of ideal for height) <b>plus</b> any TWO of following risk factors: family history of type 2 diabetes in first- or second-degree relative, member of high-risk ethnic population, maternal history of diabetes or gestational diabetes during the child's gestation, or signs of insulin resistance or of a clinical condition associated with insulin resistance, including small-for-gestational-age birth weight. FPG is the preferred test.</li> </ul>

Reference: "Standards of Medical Care in Diabetes," DIABETES CARE, Volume 32, Supplement 1, January 2009  
American Diabetes Association

QIC (03/17/09)

<p><i>Screening Criteria for Pre-Diabetes or Diagnosis of Diabetes</i> Fasting/Random plasma glucose <i>[Diagnosis must be confirmed for diagnosis of diabetes]</i></p>	<p>As indicated/annually for those with pre-diabetes</p>	<p>Note: To test for pre-diabetes or diabetes, either a FPG test or a 2-hr oral glucose tolerance test (OGTT; 75g. glucose load), or both, is appropriate.</p> <ul style="list-style-type: none"> <li>• Symptoms of diabetes and a casual plasma glucose <math>\geq 200</math> mg/dl (11.1 mmol/l).</li> <li>• Fasting plasma glucose (FPG) <math>\geq 126</math> mg/dl (7.0 mmol/l). Fasting is defined as no caloric intake for at least 8 hours.</li> <li>• 2-hr plasma glucose (PG) <math>\geq 200</math> mg/dl (11.1 mmol/l) during an oral glucose tolerance test (OGTT). The test should be performed as described by the World Health Organization (WHO), using a glucose load containing the equivalent of 75-g anhydrous glucose dissolved in water.</li> </ul>																		
<p>Diagnosis of Gestational DM with a 100-g glucose load</p>	<p>As indicated/Note: Women with GDM should be screened for diabetes 6-12 weeks postpartum and should be followed up with subsequent screening for the development of diabetes or pre-diabetes.</p>	<table border="1"> <thead> <tr> <th></th> <th>mg/dl</th> <th>Mmol/l</th> </tr> </thead> <tbody> <tr> <td>100-g Glucose load</td> <td></td> <td></td> </tr> <tr> <td>Fasting</td> <td><math>\geq 95</math></td> <td>5.3</td> </tr> <tr> <td>1-hr</td> <td><math>\geq 180</math></td> <td>10.0</td> </tr> <tr> <td>2-hr</td> <td><math>\geq 155</math></td> <td>8.3</td> </tr> <tr> <td>3-hr</td> <td><math>\geq 140</math></td> <td>7.8</td> </tr> </tbody> </table>		mg/dl	Mmol/l	100-g Glucose load			Fasting	$\geq 95$	5.3	1-hr	$\geq 180$	10.0	2-hr	$\geq 155$	8.3	3-hr	$\geq 140$	7.8
	mg/dl	Mmol/l																		
100-g Glucose load																				
Fasting	$\geq 95$	5.3																		
1-hr	$\geq 180$	10.0																		
2-hr	$\geq 155$	8.3																		
3-hr	$\geq 140$	7.8																		
<p>HbA1c</p>	<p>Every 3-6 months</p>	<ul style="list-style-type: none"> <li>• Goal for patients in general is <math>&lt; 7.0\%</math> (referenced to a nondiabetic range of 4.0-6.0% using DCCT-based assay)</li> <li>• Goal for the individual patient is as close to normal (<math>&lt; 6.0\%</math>) as possible without significant hypoglycemia.</li> </ul> <p>Note: more stringent control (<math>&lt; 6\%</math>) may further reduce complications at increased risk of hypoglycemia; less stringent treatment goals may be appropriate for patients with severe or frequent hypoglycemia, patients with limited life expectancies, very young children or older adults, and individuals with comorbid conditions.</p>																		
<p>FPG and OGTT values (Reference Guide)</p>		<ul style="list-style-type: none"> <li>• FPG <math>&lt; 100</math> mg/dl (5.6 mmol/l) = normal fasting glucose</li> <li>• FPG <math>\geq 100</math> mg/dl (5.6 mmol/l) and <math>&lt; 126</math> mg/dl (6.9 mmol/l) = impaired fasting glucose (IFG)</li> <li>• 2-hr plasma glucose <math>140</math> mg/dl (7.8 mmol/l) to <math>199</math> mg/dl (11.0 mmol/l) = impaired glucose tolerance (IGT)</li> </ul> <p>Note: IFG and IGT have been officially termed “pre-diabetes.” Both categories are risk factors for future diabetes and cardiovascular disease.</p>																		
<p>Fasting lipid profile</p>	<p>Annually</p>	<p>Treatment goals are: LDL <math>&lt; 100</math> mg/dl (2.6 mmol/l); triglycerides <math>&lt; 150</math> mg/dl (1.7 mmol/l) and HDL <math>&gt; 40</math> mg/dl (1.1 mmol/l) [for women, increase HDL goal by 10 mg/dl].</p> <ul style="list-style-type: none"> <li>• In individuals with diabetes aged <math>&gt; 40</math> years with total cholesterol <math>\geq 135</math> mg/dl, <i>without</i> overt cardiovascular disease, statin therapy to achieve an LDL reduction of 30-40% regardless of baseline level.</li> <li>• For persons with diabetes <math>\leq 40</math> years, <i>without</i> overt cardiovascular disease, but at increased risk due to other factors or long duration of diabetes, the addition of pharmacological therapy is appropriate.</li> <li>• Those with diabetes <i>and</i> overt cardiovascular disease are at very high risk and should be treated with a statin drug. A lower LDL goal of <math>&lt; 70</math> mg/dl (1.8 mmol/l), using a high dose of a statin, is an option in these patients.</li> </ul>																		

Reference: “Standards of Medical Care in Diabetes,” DIABETES CARE, Volume 32, Supplement 1, January 2009  
American Diabetes Association

QIC (03/17/09)

	Urinalysis	Annually	
	Urine microalbumin/creatinine	Annually	Measurement of albumin to creatinine ratio in random collection, or 24 hour with creatinine (simultaneous creatinine clearance), or timed, e.g., 4 hrs or overnight.
	Serum creatinine	Initial/annually	The serum creatinine alone should not be used as a measure of kidney function but rather used to estimate GFR and stage the level of chronic kidney disease.
	EKG	Initial	If patient is $\geq 40$ yrs old or DM $\geq 10$ years or clinically indicated
	Thyroid assessment	Initial/as indicated	Thyroid palpation, thyroid functions test(s) if indicated
<b>Management (diabetics and pre-diabetics, as applicable)</b>	Review self-management skills	Initial/ongoing	Check self-monitoring log book, diet, exercise, and meds. Refer for diabetes self-management training if indicated
	Establish/Review nutrition plan	Initial/ongoing	Refer to medical nutrition therapy if indicated. <ul style="list-style-type: none"> <li>Those with impaired glucose tolerance or impaired fasting glucose should be referred for a program aimed at reducing weight by 5-10%.</li> <li>Bariatric surgery should be considered for adults with BMI <math>\geq 35</math> kg/m<sup>2</sup> and type 2 diabetes, especially if the diabetes is difficult to control with lifestyle and pharmacologic therapy. Note: Long-term benefits, cost-effectiveness, and risks of bariatric surgery in individuals with with type 2 diabetes have not been studied in well-designed randomized clinical trials with medical and lifestyle therapy as the comparator.</li> <li>Reduction of protein intake to 0.8-1.0 g/kg body weight in individuals with diabetes and the earlier stages of chronic kidney disease (CKD) and to 0.8 g/kg body weight per day in later stages of CKD may improve measures of renal function.</li> </ul>
	Review physical activity plan	Initial/ongoing	Access/prescribe based on patient's health exams <ul style="list-style-type: none"> <li>At least 150 min/week of moderate-intensity aerobic physical activity (50-70% of maximum heart rate) and/or at least 90 min/week of vigorous aerobic exercise (&gt;70% of maximum heart rate).</li> </ul>
	Prevention/delay of type 2 diabetes:	Initial/ongoing	<ul style="list-style-type: none"> <li>In addition to lifestyle counseling, metformin may be considered in those who are at very high risk (combined impaired fasting glucose and impaired glucose tolerance plu other risk factors) and who are obese and under 60 years of age.</li> </ul>
	Immunizations <ul style="list-style-type: none"> <li>Influenza Vaccine</li> <li>Pneumococcal Vaccine</li> </ul>	Annually Recommended	<ul style="list-style-type: none"> <li>Provide at least one lifetime pneumococcal vaccine for adults with diabetes. A one-time revaccination recommended for individuals <math>\geq 65</math> years of age previously immunized when &lt;65 if administered <math>\geq 5</math> years ago.</li> </ul>
	Counseling <ul style="list-style-type: none"> <li>Tobacco use</li> <li>Psychosocial adjustment</li> <li>Sexuality/impotence</li> <li>Preconception/pregnancy</li> <li>Aspirin therapy</li> </ul>	Annually/ongoing Annually/ongoing  Ongoing, as indicated	Smoking cessation counseling  Use aspirin therapy (75-162 mg/day) or alternative antiplatelet agent for those at increased cardiovascular risk.

Reference: "Standards of Medical Care in Diabetes," DIABETES CARE, Volume 32, Supplement 1, January 2009  
American Diabetes Association

QIC (03/17/09)